



TITLE:

Buchsbaum Rings and Cone Singularities

AUTHOR(S):

MIYAZAKI, Chikashi

CITATION:

MIYAZAKI, Chikashi. Buchsbaum Rings and Cone Singularities. 数理解析
研究所講究録 1987, 621: 60-72

ISSUE DATE:

1987-04

URL:

<http://hdl.handle.net/2433/99898>

RIGHT:

Buchsbaum Rings and Cone Singularities

宮崎 哲 (Chikashi MIYAZAKI) (早大・理工)

Let $S=k(X_0, \dots, X_N)$ be the polynomial ring. Let us regard the polynomial ring S as a graded ring in which the degree of X_j is one for every $0 \leq j \leq N$. Let $\mathfrak{m} = \bigoplus_{d \geq 1} S_d$. Let M be a finitely generated graded S -module of dimension $n+1$. Then M is called a Buchsbaum S -module (resp. a quasi-Buchsbaum S -module) if $M_{\mathfrak{m}}$ is a Buchsbaum $S_{\mathfrak{m}}$ -module (resp. a quasi-Buchsbaum $S_{\mathfrak{m}}$ -module). That is, M is a quasi-Buchsbaum module if and only if $\mathfrak{m}H_{\mathfrak{m}}^i(M) = 0$ for any $0 \leq i \leq n$. On the other hand, M is a Buchsbaum module if and only if the difference $l(M/qM) - e_q(M)$ is an invariant for any homogeneous parameter ideal $q (\subset \mathfrak{m})$ for M , where l , respectively e , denotes length, respectively multiplicity of q .

In this paper, we search for the condition that a graded S -module M is a Buchsbaum module. In particular, our purpose is to clarify the difference between Buchsbaum property and quasi-Buchsbaum property.

For this, we introduce the notion of quasi-Buchsbaum modules of type r in Section 1. Then we give a criterion (Theorem 1.7) for it, which explicitly describes "Surjectivity Criterion" through spectral sequence. In Corollary 1.10, we give a sufficient condition that M is Buchsbaum. It is described

only by the vanishing of its local cohomologies. This generalizes a result of Goto-Watanabe (4, Proposition 3.1). Also, the theorem enables us to calculate the length of $\text{Ext}_S^{n+1}(k, M)$ in Corollary 1.12.

In Section 2, we investigate the divisors of Segre products. In particular, we construct very concrete examples of quasi-Buchsbaum but not Buchsbaum rings.

§1. Quasi-Buchsbaum modules of type r

Definition 1.1. M is a quasi-Buchsbaum module of type r if, for every homogeneous system f_1, \dots, f_{r-1} of parameters, $M/(f_1, \dots, f_k)M$ is a quasi-Buchsbaum module for every $k \leq r-1$.

Remark 1.2. In the above definition, we have only to take elements of degree one as homogeneous systems of parameters, by Goto-Suzuki (6, Theorem 1.9) or Suzuki (19, Theorem 3.6).

Remark 1.3. M is a quasi-Buchsbaum module of type $n+1$ if and only if M is a Buchsbaum module, by Stückrad-Vogel (14).

Let U_i be an open set $D_+(X_i)$ in $\text{Proj } S$ for $0 \leq i \leq N$. Then $\mathcal{U} = \{U_i\}_{0 \leq i \leq N}$ is an open covering of $\text{Proj } S$. Let us consider a Čech complex $C^\bullet = C^\bullet(\mathcal{U}; \tilde{M})$, where \tilde{M} is the sheaf associated to the graded S -module M on $\text{Proj } S$. Then we define a complex

$L' = (0 \longrightarrow M \xrightarrow{\varepsilon} C'(-1))$, where ε is a natural map. (e.g. See Godement (2).)

Let K be the Koszul complex associated to $\langle X_0, \dots, X_N \rangle$, that is, $K_p = \bigwedge^p \left(\bigoplus_{k=0}^N Ae_k \right) = \bigoplus_{0 \leq i_1 < \dots < i_p \leq N} Ae_{i_1} \wedge \dots \wedge e_{i_p}$, where $\langle e_0, \dots, e_N \rangle$ is a free basis.

In this way, we have a double complex $K' = \text{Hom}(K, L')$. Now let us consider two "stupid" filtrations, that is,

$$K'_t = \sum_{p \geq t} K^{p,q} \quad \text{and} \quad K'_t = \sum_{q \geq t} K^{p,q}.$$

As usual, we get the spectral sequence for each filtered complex:

$$\begin{aligned} E_1^{p,q} &= \text{Ker } d^{-p,q} / \text{Im } d^{-p,q-1} \\ E_1^{p,q} &= \text{Ker } d^{-p,q} / \text{Im } d^{-p-1,q} \end{aligned} \quad \begin{aligned} &\Rightarrow \\ &\Rightarrow \end{aligned} \quad H^{p+q}(K' \cdot \cdot)$$

Lemma 1.4. Under the above conditions, we have the following things.

$$(1) \quad E_1^{p,q} = \bigwedge^p \left(\bigoplus_{k=0}^N H_m^q(M) e_k^* \right),$$

where $\langle e_0^*, \dots, e_N^* \rangle$ is the dual basis of $\langle e_0, \dots, e_N \rangle$, and

$$(2) \quad E_1^{p,q} = \begin{cases} \text{Ext}_S^p(k, M) & \text{if } q=0 \\ 0 & \text{otherwise.} \end{cases}$$

This implies that $H^{p+q}(K' \cdot \cdot) = \text{Ext}_S^{p+q}(k, M)$.

From now on we will treat the first filtration. So we

write $E_r^{p,q}$ for $E_r^{p,q}$.

Lemma 1.5. The spectral sequence $(E_r^{p,q})$ does not depend on the choice of coordinates $\langle X_0, \dots, X_N \rangle$.

Now then we will study the relation between the spectral sequence and Buchsbaum property.

Proposition 1.6. Notations being above, the following conditions are equivalent.

- (a) M is a quasi-Buchsbaum S -module.
- (b) $d_1^{p,q} : E_1^{p,q} \longrightarrow E_1^{p+1,q}$ is a zero map for $q \leq n$.
- (c) $d_1^{0,q} : E_1^{0,q} \longrightarrow E_1^{1,q}$ is a zero map for $q \leq n$.

Proof. By (1.4.1.), $E_1^{p,q} = \wedge^p \left(\bigoplus_{k=0}^N H_m^q(M) e_k^* \right)$. From the construction of the double complex $K^{\cdot,\cdot}$, we see that

$$d_1^{p,q}(e_{i_1}^* \wedge \dots \wedge e_{i_p}^*) = \sum_{j=0}^N X_j e_j^* \wedge e_{i_1}^* \wedge \dots \wedge e_{i_p}^*.$$

Thus the assertion follows from the definition of quasi-Buchsbaum module.

Suppose that M is a quasi-Buchsbaum module and that every subset I of $\{X_0, \dots, X_N\}$ such that $\#I = n+1$ makes a system of parameters of M . Setting $\bar{M} = M/X_j M$, we have the following exact sequence:

$$0 \longrightarrow (0:X_j)_M \longrightarrow M \xrightarrow{\cdot X_j} M \longrightarrow \bar{M} \longrightarrow 0.$$

Since $H_m^q((0:X_j)_M) = 0$ for $q \geq 1$ and M is a quasi-Buchsbaum module, we have the following short exact sequence:

$$0 \longrightarrow H_m^{q-1}(M) \longrightarrow H_m^{q-1}(\bar{M}) \longrightarrow H_m^q(M) \longrightarrow 0.$$

for $1 \leq q \leq n$. Thus we have the following commutative diagram with exact rows:

$$(1.7.1) \quad \begin{array}{ccccccc} 0 & \longrightarrow & H_m^{q-1}(M) & \longrightarrow & H_m^{q-1}(\bar{M}) & \xrightarrow{\alpha} & H_m^q(M) \longrightarrow 0 \\ & & \downarrow \phi & & \downarrow \bar{\psi} & & \downarrow \phi' \\ 0 & \longrightarrow & H_m^{q-1}(M) & \xrightarrow{\beta} & H_m^{q-1}(\bar{M}) & \longrightarrow & H_m^q(M) \longrightarrow 0 \end{array}$$

for $1 \leq q \leq n$, where ϕ , ϕ' and $\bar{\psi}$ are multiplication maps $\cdot X_i$'s. Since M is a quasi-Buchsbaum module, ϕ and ϕ' are zero maps. By snake lemma, we get a graded S -homomorphism of degree 2 $\psi : H_m^q(M) \longrightarrow H_m^{q-1}(M)$ such that $\psi \cdot \alpha = \beta \cdot \bar{\psi}$. Let us write $(X_i \wedge X_j)$ for this map ψ . Since $\psi = 0$ is equivalent $\bar{\psi} = 0$, we see that \bar{M} is a quasi-Buchsbaum module if and only if $(X_i \wedge X_j)$ is a zero map for any i . In other words, M is a quasi-Buchsbaum module of type 2 if and only if and only if $(X_i \wedge X_j)$ are zero map for any i and j .

Continuing the similar steps, we have the following theorem.

Theorem 1.7. Let S be the polynomial ring $k(X_0, \dots, X_N)$.

Let M be a finitely generated graded S -module with dimension $n+1$. Suppose that M is a quasi-Buchsbaum module of type $r-1$. Then

$$(X_I) = (X_{i_1} \wedge \dots \wedge X_{i_r}) : H_m^q(M) \longrightarrow H_m^{q-r+1}(M)$$

is well-defined.

Furthermore, M is a quasi-Buchsbaum module of type r if and only if (X_I) defined above is a zero map for any I .

Now there still remains to link between the spectral sequence $\{E_r^{p,q}\}$ and the maps (X_I) 's constructed above.

Lemma 1.8. Let M be a quasi-Buchsbaum graded S -module of type $r-1$ and with dimension $n+1$. Then we have

$$(1) \quad E_r^{p,q} = \bigwedge^p \left(\bigoplus_{k=0}^N H_m^q(M) e_k^* \right) \quad \text{for any } q \neq n+1.$$

By (1), we can write $(d_r^{p,q})_{J,K} : H_m^q(M) \longrightarrow H_m^{q-r+1}(M)$ for the map from e_K^* -component to $e_J \wedge e_K^*$ -component of the map $d_r^{p,q}$.

Then we have

$$(2) \quad (d_r^{p,q})_{J,K} = (-1)^{(r-1)(p+q-r)} (X_J), \quad \text{where } (X_J) \text{ is the map defined in (1.7).}$$

Then we have the following theorem.

Theorem 1.9. Notations being above, the following conditions are equivalent.

- (a) M is a quasi-Buchsbaum S -module of type r .

(b) $d_s^{p,q} : E_s^{p,q} \longrightarrow E_s^{p+s, q-s+1}$ is a zero map for $s \leq r$ and $q \leq n$.

(c) $d_s^{0,q} : E_s^{0,q} \longrightarrow E_s^{s, q-s+1}$ is a zero map for $s \leq r$ and $q \leq n$.

Corollary 1.10. Let M be a finitely generated graded S -module. Let us define

$$\mathcal{G} = \{(i, \ell) \mid 0 \leq i \leq n, \ell \in \mathbb{Z}, H_m^i(M)_\ell \neq 0\}.$$

Suppose that \mathcal{G} satisfies the following condition (*).

(*) For any (i, ℓ) and (j, m) of \mathcal{G} , if $i \geq j$, then $i + \ell + 1 \neq j + m$.

Then M is a Buchsbaum module.

Proof. Since $(E_1^{p,q})_\ell = \oplus H_m^q(M)_\ell$, $(E_1^{p+s, q-s+1})_{\ell+s} = \oplus H_m^{q-s+1}(M)_{\ell+s}$ and $d_s^{p,q}$ is a graded homomorphism of degree s , the assumption gives that $d_s^{p,q}$ is zero for any $p, q (\leq n)$ and s . By Theorem 1.9 (or (1.7)), M is a Buchsbaum module.

Remark 1.11. (1.10) is a generalization of Goto-Watanabe (4) and Schenzel (13, Theorem 3.1) and is "best possible" in the following sense. Unless \mathcal{G} satisfies (*), we cannot see only by the vanishing of its local cohomologies whether or not M is a Buchsbaum module. In fact, Goto abstractly and systematically constructed such examples in his papers (3, 5). Moreover, his method gives the construction

of quasi-Buchsbaum ring of type r but not of type $r+1$ by virtue of Evans-Griffith (1).

Now then, let us calculate the length of $\text{Ext}_S^{n+1}(k, M)$ written by $r(M)$. By (1.9), we have the following result.

Corollary 1.12. Suppose that M is a Buchsbaum S -algebra, we have

$$\sum_{j=0}^n \binom{N+1}{n+1-j} \dim H_m^j(M) + 1 \leq r(M) \leq \sum_{j=0}^n \binom{N+1}{n+1-j} \dim H_m^j(M) + \mu(K_M),$$

where $K_M = \text{Ext}_S^{N-n}(M, S)$ and $\mu(K_M)$ is the minimal number of generators of K_M .

§2. Divisors on Segre products

Let k be a field. Let X be an arithmetically Cohen-Macaulay subscheme of $\mathbb{P}_k^N = \text{Proj } S$, that is, its affine cone $C(X) = \text{Spec } S/J$ is locally Cohen-Macaulay, where $J = \bigoplus_{t \in \mathbb{Z}} r(\mathcal{I}_{X/P}(t))$.

Let V be a subscheme of X such that $0 < \dim V = n < \dim X$. Let A be the coordinate ring of V . The following proposition shows that the results of Section 1 can be applied to geometric case.

Proposition 2.1. Under the above conditions, we have

$$\tau_0^{n+1} \text{R}\Gamma\left(\bigoplus_{t \in \mathbb{Z}} \mathcal{I}_{V/X}(t)\right) \simeq \tau_0^{n+1} \text{R}\Gamma_m(A)$$

in the derived category $D_h^+(S)$ of complexes bounded below of

graded S -modules.

Example 2.2. Let $X = \mathbb{P}_k^r \times \mathbb{P}_k^s$ be Segre embedding in $\mathbb{P} = \mathbb{P}_k^{rs+r+s}$. Let V be a divisor of X corresponding to $\mathcal{O}_X(a, b) = p_1^* \mathcal{O}_{\mathbb{P}_k^r}(a) \otimes p_2^* \mathcal{O}_{\mathbb{P}_k^s}(b)$.

(1) V is an arithmetically Cohen-Macaulay subscheme of \mathbb{P} if and only if $a-r \leq b \leq a+s$.

(2) The following things are equivalent.

(a) V is an arithmetically Buchsbaum subscheme of \mathbb{P} .

(b) V is an arithmetically quasi-Buchsbaum subscheme of \mathbb{P} .

(c) $a-r-1 \leq b \leq a+s+1$.

Proof. It is given, for example, by Goto-Watanabe (7), Stückrad-Vogel (18) or Schenzel (13, Proposition 5.1). We will prove it, however, because our proof indicates the motivation of the next example.

Now let us assume $a \geq b$. First of all, let us find the numbers $1 \leq i < r+s$ and $\ell \in \mathbb{Z}$ satisfying $H^i(\mathcal{I}_{V/X}(\ell)) = 0$. By the way $\mathcal{I}_{V/X}$ is isomorphic to $\mathcal{O}_X(-a, -b)$. By Künneth formula, we have $H^i(\mathcal{I}_{V/X}(\ell)) = 0$ if and only if $i=r$ and $b \leq \ell \leq a-r-1$. This shows (1) and "(c) \Rightarrow (a)" of (2). It is trivial that (a) implies (b). Thus it still remains to prove that if $a-r-1 > b$, then V is not an arithmetically quasi-Buchsbaum subscheme of \mathbb{P} .

Let us take $\mathbb{P}^r \times \mathbb{P}^s = \text{Proj } k(X_0, \dots, X_r) \times \text{Proj } k(Y_0, \dots, Y_s)$.

Then we can show

$$\cdot X_j: H^r(\mathcal{O}_{\mathbb{P}^r}(\ell-a)) \longrightarrow H^r(\mathcal{O}_{\mathbb{P}^r}(\ell+1-a))$$

is surjective and

$$\cdot Y_j: H^0(\mathcal{O}_{\mathbb{P}^s}(\ell-b)) \longrightarrow H^0(\mathcal{O}_{\mathbb{P}^s}(\ell+1-b))$$

is injective. On the other hand we see

$$H^r(\mathcal{I}_{V/X}(\ell)) \cong H^r(\mathcal{O}_{\mathbb{P}^r}(\ell-a)) \oplus H^0(\mathcal{O}_{\mathbb{P}^s}(\ell-b)).$$

Hence we have that $\bigoplus_{\ell \in \mathbb{Z}} H^i(\mathcal{I}_{V/X}(\ell))$ is not a k -vector space if

$a-r-1 > b$. This gives V is not an arithmetically Buchsbaum subscheme of \mathbb{P} .

In Example 2.2, we need not distinguish Buchsbaum rings from quasi-Buchsbaum rings. However, taking $X = \mathbb{P}_k^r \times \mathbb{P}_k^r \times \mathbb{P}_k^r$, we can find the examples clarifying the difference between them.

Theorem 2.3. Let $X = \mathbb{P}_k^r \times \mathbb{P}_k^r \times \mathbb{P}_k^r$ be Segre embedding in $\mathbb{P} = \mathbb{P}_k^{(r+1)^3-1}$. Let V be a divisor of X corresponding to $\mathcal{O}_X(a-r-1, a, a+r+1)$. Then V is an arithmetically quasi-Buchsbaum subscheme of type r in \mathbb{P} but not an arithmetically quasi-Buchsbaum subscheme of type $r+1$.

Proof. For simplicity, we will prove only in case $r=1$. Let $\langle X, Y \rangle \times \langle Z, W \rangle \times \langle U, T \rangle$ be a coordinate of $X = \mathbb{P}_k^1 \times \mathbb{P}_k^1 \times \mathbb{P}_k^1$. By Künneth formula, we have

$$H^2\left(\bigoplus_{\ell \in \mathbb{Z}} \mathcal{F}_{V/X}(\ell)\right) = k \cdot 1 \otimes k \cdot \frac{1}{Z \cdot W} \otimes \left(k \cdot \frac{1}{U^3 T} + k \cdot \frac{1}{U^2 T^2} + k \cdot \frac{1}{U T^3}\right)$$

and

$$H^1\left(\bigoplus_{\ell \in \mathbb{Z}} \mathcal{F}_{V/X}(\ell)\right) = (k \cdot X^2 + k \cdot XY + k \cdot Y^2) \otimes k \cdot 1 \otimes k \cdot \frac{1}{U \cdot T}.$$

Then we see

$$(X \otimes Z \otimes U) \wedge (Y \otimes W \otimes T) : H^2\left(\bigoplus_{\ell \in \mathbb{Z}} \mathcal{F}_{V/X}(\ell)\right) \longrightarrow H^1\left(\bigoplus_{\ell \in \mathbb{Z}} \mathcal{F}_{V/X}(\ell)\right)$$

is not zero. By Theorem 1.7, V is not arithmetically quasi-Buchsbaum of type 2. Further, it is easy to show V is arithmetically quasi-Buchsbaum. Thus the assertion is proved.

References

1. E. G. Evans and P. Griffith, Local cohomology modules for normal domains, J. London Math. Soc. 19(1979), 277-284.
2. R. Godement, Topologie algébrique et théorie des faisceaux, Hermann, Paris (1958).
3. S. Goto, On Buchsbaum rings, J. Algebra, 67(1980), 272-279.
4. S. Goto, Buchsbaum rings of maximal embedding dimension, J. Algebra, 76(1982), 383-399.
5. S. Goto, A note on quasi-Buchsbaum rings, Proc. Amer. Math. Soc. 90(1984) 511-516.
6. S. Goto and N. Suzuki, Index of reducibility of parameter ideals in a local ring, J. Algebra, 87(1984), 53-88.
7. S. Goto and K.-i. Watanabe, On graded rings I, J. Math. Soc.

Japan, 30(1978), 179-213.

8. A. Grothendieck, Local cohomology, Lecture Note in Math. 41, Springer-Verlag, Heidelberg(1967).
9. R. Hartshorne, Residue and duality, Lecture Note in Math. 20, Springer-Verlag, Heidelberg(1966).
10. M. Hochster, Topics in the homological theory of modules over commutative rings, Regional Conference Series in Math. 24, Amer. Math. Soc., Providence, 1975.
11. M. Hochster, Canonical elements in local cohomology modules and the direct summand conjecture, J. Algebra, 84(1983), 503-553.
12. M. Ishida, On the dualizing complexes of normal isolated Du Bois singularities, Preprint.
13. P. Schenzel, Application of dualizing complexes to Buchsbaum rings, Advances in Math. 44(1982), 61-77.
14. P. Schenzel, Dualisierende Komplexe in der lokalen Algebra und Buchsbaum-Ringe, Lecture Note in Math. 907, Springer-Verlag, Heidelberg(1980).
15. J. Stückrad, Über die kohomologische Charakterisierung von Buchsbaum-Moduln, Math. Nachr. 95(1980), 265-272.
16. J. Stückrad and W. Vogel, Eine Verallgemeinerung der Cohen-Macaulay-Ringe und Anwendungen auf ein Problem der Multiplizitätstheorie, J. Math. Kyoto Univ. 13(1973), 513-528.
17. J. Stückrad and W. Vogel, Toward a theory of Buchsbaum singularities, Amer. J. Math. 100(1978), 727-746.
18. J. Stückrad and W. Vogel, On Segré product and

applications, J. Algebra, 54(1978), 374-389.

19. J. Stückrad, P. Schenzel and W. Vogel, Theorie der Buchsbaum-Moduln, Preprint der Sektion Mathematik, No.23/24 Martin-Luther-Universität, Halle, 1979.

20. N. Suzuki, On quasi-Buchsbaum modules — an application of theory of FLC-modules, Preprint.

21. N. V. Trung, Toward a theory of generalized Cohen-Macaulay modules, Nagoya Math. J, 102(1986), 1-49.

22. W. Vogel, A non-zero-divisor characterization of Buchsbaum modules, Michigan Math. J. 28(1981), 147-152.

23. K.-i. Watanabe, Some remarks concerning Demazure's construction of normal graded rings, Nagoya Math. J. 83(1981), 203-211.

Department of Mathematics
School of Science and Engineering
Waseda University
3-4-1, Okubo, Shinjuku-ku,
Tokyo, 160 Japan